

Luminosity segregation versus fractal scaling in the galaxy distribution

MARTIN KERSCHER¹

¹ *Sektion Physik, Ludwig-Maximilians-Universität, Theresienstraße 37, D-80333 München, Germany*

PACS. 98.65.-r – Galaxy groups, clusters, and superclusters; large scale structure of the Universe.

PACS. 98.62.Ve – Statistical and correlative studies of properties (luminosity and mass functions; mass-to-light ratio; Tully- Fisher relation, etc.) .

PACS. 89.75.Da – Systems obeying scaling laws.

Abstract. – In this letter I present results from a correlation analysis of three galaxy redshift catalogs: the SSRS2, the CfA2 and the PSCz. I will focus on the observation that the amplitude of the two-point correlation function rises if the depth of the sample is increased. There are two competing explanations for this observation, one in terms of a fractal scaling, the other based on luminosity segregation. I will show that there is strong evidence that the observed growth is due to a luminosity dependent clustering of the galaxies.

Introduction. – One of the problems in cosmology is to understand the formation of the large-scale structures in the Universe, as traced by the spatial distribution of galaxies. Theoretical models of large-scale structure and galaxy formation, whether involving analytical predictions or numerical simulations, are based on some form of random or stochastic initial conditions. This means that a statistical interpretation of the observed galaxy distribution is required, and that statistical tools must be deployed in order to discriminate between different cosmological models (e.g. [1]). The most frequently employed statistical measure is the two-point correlation function. Higher-order correlations are important, but already the observed two-point correlation properties of the galaxy distribution impose strong constraints on the models of structure formation. However, two basically different interpretations of the observed two-point properties are discussed: in the “standard” picture the galaxy distribution is assumed to be homogeneous on large scales. The correlations of the deviations from this homogeneous distribution are quantified by the two-point correlation function $\xi(r)$ (see e.g. [2]). In the alternative picture the galaxy distribution is modelled as a fractal. The two-point correlations are then measured with the conditional density $\Gamma(r)$ (see e.g. [3]). The inhomogeneous nature of a fractal challenges the standard picture. Clearly, these two models lead to different interpretations of observational results. In this letter I will focus on the growing of the amplitude of the two-point correlation function $\xi(r)$ with the sample depth. This growing amplitude is either explained with the scaling properties of a fractal or with luminosity segregation, a luminosity dependent clustering strength. By reanalysing three galaxy catalogues I will show that there are strong arguments in favour of luminosity segregation.




Fig. 1




Fig. 2

Fig. 1 – Luminosity L of the galaxies in the Southern Sky Redshift Survey 2 (SSRS2) plotted against their distance s to our galaxy. Volume-limited samples comprise the galaxies in the upper left part enclosed by the solid lines ($120h^{-1}\text{Mpc}$), dotted lines ($100h^{-1}\text{Mpc}$), short dashed lines ($90h^{-1}\text{Mpc}$), long dashed lines ($80h^{-1}\text{Mpc}$), and dashed-dotted lines ($70h^{-1}\text{Mpc}$). Low luminosity galaxies at large distances are not observed, as can be seen from the empty region in the lower right part.

Fig. 2 – Sample geometry of a volume-limited sample with sample depth R (simplified sketch). Only galaxies inside the opening angle Ω and with a distance $s < R$ enter the sample.

Two-point correlations. – Let me first discuss the stochastic picture where the galaxies positions in space are treated as a realization of a random process. The product density $\rho_2(\mathbf{x}_1, \mathbf{x}_2)dV(\mathbf{x}_1)dV(\mathbf{x}_2)$ is the probability of finding two galaxies in the volume elements $dV(\mathbf{x}_1)$ and $dV(\mathbf{x}_2)$, respectively. In a homogeneous and isotropic point distribution with mean number density ρ one defines the two-point correlation function $\xi(r)$ (e.g. [2])

$$\rho_2(\mathbf{x}_1, \mathbf{x}_2) = \rho^2(\xi(r) + 1), \quad (1)$$

with $r = |\mathbf{x}_1 - \mathbf{x}_2|$. The conditional density can be defined as $\Gamma(r) \equiv \rho(\xi(r) + 1)$. For a point distribution on a fractal the mean number density ρ , and also $\xi(r)$ are not well defined. Thus, in this case, one investigates the two-point correlations with the conditional density $\Gamma(r)$: the density of galaxies at a distance of r as seen from another galaxy [4]. A scale invariant cumulant $\xi(r)$ is typically found in critical systems, whereas a scale invariant conditional density $\Gamma(r)$ is an indication for a fractal system. Clearly, only in the limit $r \rightarrow 0$, both $\xi(r)$ and $\Gamma(r)$ may show the same scaling behaviour. An instructive discussion of the different scaling regimes in the galaxy distribution is presented by [5].

Galaxy samples. – In a typical galaxy catalogue the position on the sky inside a given angular region Ω and the flux in a given wave-band are measured. The distance to our position is estimated from the redshift of the galaxy. A flux-limited sample consists out of galaxies down to a limiting flux f_{lim} . To study the clustering properties of such a galaxy catalogue, I extract a sequence of volume-limited samples. A volume-limited subsample is constructed by introducing a limiting depth R and a limiting luminosity L_{lim} and by admitting only galaxies within a distance $s \leq R$ from our position and a luminosity $L \geq L_{\text{lim}}$ ($L_{\text{lim}} \propto R^2 f_{\text{lim}}$, in the Euclidean case, see Fig. 1). For a given sample one defines the sample-dependent number density $\rho_S = N/V$, with the number of galaxies N , and the volume V . The two-point correlation function ξ_S , and the conditional density Γ_S determined from this sample satisfy the relation $\Gamma_S(r) = \rho_S(1 + \xi_S(r))$. If our galaxy sample stems from a homogeneous distribution, neither ξ_S nor Γ_S should change if one increases the sample size (despite statistical fluctuations). If




Fig. 3 – $\xi(r)$ and $\Gamma(r)$ for volume limited samples from the SSRS2 with depth $120h^{-1}\text{Mpc}$ (solid line), $100h^{-1}\text{Mpc}$ (dotted line), $90h^{-1}\text{Mpc}$ (short dashed line), $80h^{-1}\text{Mpc}$ (long dashed line), and $70h^{-1}\text{Mpc}$ (dashed-dotted line). The limiting luminosity changes in these samples. The labels *LS* and *minus* mark the results obtained with the Landy & Szalay and the minus estimator, respectively. The solid dot marks $r_0(120h^{-1}\text{Mpc})$, the open dots mark $r_0(100h^{-1}\text{Mpc})$, $r_0(90h^{-1}\text{Mpc})$, $r_0(80h^{-1}\text{Mpc})$, and $r_0(70h^{-1}\text{Mpc})$ from right to left, according to Eq.(2). The smooth solid line is $\xi(r)$ for a fractal with $D = 2$ and $r_0(70h^{-1}\text{Mpc})$ according to Eq.(2). Two- σ error bars, determined from a Poisson process, are shown only for the $120h^{-1}\text{Mpc}$ and $70h^{-1}\text{Mpc}$ samples.

our sample stems from a point distribution on a fractal the number density ρ_S is depending on the sample size, but the scaling exponent $D - 3$ of $\Gamma_S(r) \propto r^{D-3}$ stays invariant (D is the correlation dimension). As a result ξ_S is changing with the size of the sample (see e.g. [3]). For the two-point correlation function often the following parameterisation is used $\xi_S(r) = \left(\frac{r_0}{r}\right)^\gamma$ with the scaling index γ . The so-called “correlation length” r_0 quantifies the amplitude r_0^γ of a scale invariant correlation function. Consider a large sample with a depth $R_{\max}$ and several smaller samples $R \leq R_{\max}$ within (see Fig. 2). On a fractal r_0 is proportional to R [6], specifically

$$r_0(R) = \frac{R}{R_{\max}} r_0(R_{\max}), \text{ and } \xi(r) = 2 \left(\frac{r}{r_0(R)} \right)^{D-3} - 1. \quad (2)$$

In the following I will estimate the two-point correlation function as well as the conditional density for three galaxy samples. I checked that for the samples and the scales considered here, the estimators discussed by [7] give consistent results. I illustrate this by showing the results for $\xi(r)$ both for the minus (reduced sample) estimator as favoured by [3], and the estimator due to Landy & Szalay [8]. [7] showed that the Landy & Szalay estimator has preferable variance properties.

Luminosity segregation but no fractal scaling in the SSRS2. – The Southern Sky Redshift Survey 2 (SSRS2, [9]) is 99% complete with a limiting magnitude of $m_{\text{lim}} = 15.5$ (magnitudes are logarithmic flux measures). The angular extent is $-40^\circ \leq \delta \leq -2.5^\circ$ with $b \leq -40^\circ$ and $\delta \leq 0^\circ$ with $b \geq 35^\circ$ (declination δ , galactic latitude b). The magnitudes were K -corrected as described in [10], and luminosity distances were used. Nearly identical results could be obtained using Euclidean distances and no K -correction. In Fig. 3 both $\xi(r)$ and $\Gamma(r)$ are shown for a sequence of volume-limited samples from the SSRS2. The number density in the volume-limited samples decreases from $70h^{-1}\text{Mpc}$ to $120h^{-1}\text{Mpc}$. Consequently, the conditional density $\Gamma(r)$ is decreasing with the sample depth. The amplitude of $\xi(r)$ increases with the depth of the samples, and r_0 roughly follows the relation (2). This was interpreted as a sign of a fractal galaxy distribution (e.g. [3]). However, another explanation is possible.




Fig. 4 – $\xi(r)$ and $\Gamma(r)$ for subsamples from the volume-limited samples from the SSRS2 with a depth of $R_{\max} = 120h^{-1}\text{Mpc}$. The depth of the sub-samples is $120h^{-1}\text{Mpc}$ (solid line), $100h^{-1}\text{Mpc}$ (dotted line), $90h^{-1}\text{Mpc}$ (short dashed line), $80h^{-1}\text{Mpc}$ (long dashed line), and $70h^{-1}\text{Mpc}$ (dashed-dotted line). All samples have the same limiting luminosity. The smooth solid line is $\xi(r)$ for a fractal with $D = 2$ and $r_0(70h^{-1}\text{Mpc})$ according to Eq.(2). Marks as in Fig. 3.

Due to the construction of the volume-limited sample the mean absolute luminosity of the galaxies in the sample increases with the depth of the sample (compare with Fig. 1). Hence, the growing amplitude of the correlation function for deeper volume-limited samples may be a result of the stronger clustering of the more luminous galaxies. This is called luminosity segregation. Clearly, the two-point correlation function $\xi(r)$ applied to a sequence of volume-limited samples is not able to distinguish between both claims. To test the scaling relation (2) independent from any luminosity dependence I use a volume-limited sample with a depth of $R_{\max} = 120h^{-1}\text{Mpc}$. From this volume-limited sample I extract a sequence of subsamples with depths of $R \leq R_{\max}$ (see Fig. 2). All these subsamples have the same lower limit in luminosity (see Fig. 1). As can be seen from Fig. 4 the estimated $\xi(r)$ are consistent in these samples but inconsistent with the fractal prediction. There is no indication for a fractal scaling of the “correlation length” $r_0(R)$ as given in Eq.(2). Moreover, the conditional densities $\Gamma(r)$ of these samples nearly overlap. Measurement errors, e.g. for the position of the galaxies, have no visible effect on the correlation functions. The dominant contribution is the statistical error. However there is only one realisation of the galaxy distribution in the Universe. Therefore, I have to assume a model to quantify the statistical errors. The simplest model is a purely random distribution of points, the Poisson process. I estimate the errors from 100 realisations of a Poisson process with the sample geometry, and the number density as in the galaxy samples. Later on I will show that a more realistic modelling leads to larger errors. These errors are within the same order as determined from a Poisson process. For both models, the errors are smaller than the predicted effects from fractal scaling.

Luminosity segregation but no fractal scaling in the CfA2. – As a spatially complementary sample to the SSRS2 I use a galaxy sample from the CfA galaxy catalogue (see [11] and references therein), with $\delta \geq 0$, and $|b| \geq 35$ and a limiting magnitude of $m_{\text{lim}} = 15.5$. From this galaxy catalogue I extract similar volume-limited samples as for the SSRS2. The results shown in Figs. 5,6 lead to the same interpretation as for the SSRS2: the growing amplitude of $\xi(r)$ is caused by luminosity segregation.

Neither luminosity segregation nor fractal scaling in the PSCz. – Both the galaxies in the SSRS2 and the CfA2 were selected in the optical waveband. The galaxies in the PSCz




Fig. 5




Fig. 6

Fig. 5 – $\xi(r)$ for a sequence of volume-limited samples with changing limiting Luminosity L_{lim} from the CfA2 determined with the Landy&Szalay estimator. The smooth solid line is $\xi(r)$ for a fractal with $D = 2$ and $r_0(70h^{-1}\text{Mpc})$ according to Eq.(2). Marks as in Figs. 3.

Fig. 6 – $\xi(r)$ for samples with varying depth but with the same limiting luminosity extracted from a volume-limited sample of the CfA2 with a depth of $R_{\text{max}} = 120h^{-1}\text{Mpc}$. Marks as in Fig. 4.

survey were selected according to their flux in the infrared as detected by the IRAS satellite. A detailed description of the IRAS PSCz galaxy catalogue may be found in [12]. I extract volume-limited samples from the PSCz survey using luminosity distances within the standard masked area. I approximate the sample geometry by two spherical caps with galactic latitude $b \geq 5^\circ$ for the northern part and with $b \leq -5^\circ$ for the southern part. Hence, I neglect some regions which were left empty due to galactic absorption or confusion in the IRAS PSC maps. I filled these empty regions with random points assuming the same number density as in the fully sampled region. No differences in the correlation properties between the filled and unfilled samples are visible in the two-point measures. The $\xi(r)$ determined from a sequence of volume-limited samples is inconsistent with the fractal prediction and shows no significant variation of r_0 with the sample size (Fig. 7). As expected, extracting subsamples from one volume-limited sample with $R_{\text{max}} = 120h^{-1}\text{Mpc}$ does not change this behaviour, although the fluctuations increase in the sparser samples (Fig. 8). Clearly, there is neither an indication for a fractal scaling of r_0 with the sample depth nor for luminosity segregation (see also [13]). Due to the selection of galaxies in the infrared one does miss early type (e.g. elliptical) galaxies. Because of this selection one does not find luminosity segregation in the PSCz [14]. To go beyond the error estimates relying on the Poisson process I estimate the errors for $\xi(r)$ from the fluctuations between eleven mock galaxy catalogues, constructed from an N -body simulation based on a ΛCDM cosmology⁽¹⁾. For such a clustered point distribution, the error bars are larger than in the case of the Poisson process, but still of the same order (see Figs. 7,8). In both models, these errors are smaller than the predicted effects of fractal scaling on $\xi(r)$ (two- σ error bars are shown in the plots).

Summary. – By analysing three different galaxy catalogues I could show that the amplitude of the correlation function $\xi(r)$ is depending on the luminosity. Luminosity segregation already has been found in the SSRS2, CfA2 and also the 2dF and the SDSS galaxy catalogues (see e.g. [16,17,18,10,19,20,21]). In these investigations mainly volume-limited samples with varying depths or directly flux-limited samples have been used. Using such samples one is

⁽¹⁾A description of the procedure and references to articles describing the simulation and the construction of the mock catalogues can be found in [15].




Fig. 7




Fig. 8

Fig. 7 – $\xi(r)$ for a sequence of volume-limited samples with changing limiting luminosity L_{lim} from the PSCz determined with the Landy & Szalay estimator with marks similar to Fig. 3. The smooth solid line is $\xi(r)$ for a fractal with $D = 2.2$ and $r_0(70h^{-1}\text{Mpc})$ according to Eq.(2). Two- σ error bars are shown only for the $120h^{-1}\text{Mpc}$ sample. The larger error bars were determined from the mock catalogues. The Poisson error bars are slightly shifted to the left.

Fig. 8 – $\xi(r)$ for samples with varying depth but with the same limiting Luminosity L_{lim} extracted from a volume-limited sample of the PSCz with a depth of $R_{\text{max}} = 120h^{-1}\text{Mpc}$ similar to Fig. 4. Marks as in Fig. 7, but the two- σ error bars are shown only for the $70h^{-1}\text{Mpc}$ sample.

not able to separate the influence of luminosity segregation from a possible fractal scaling. In samples with a fixed lower limit in the luminosity and then reducing the depth of the samples I found no significant changes in the correlation function. Specifically, I found no sign for a growth of the correlation length r_0 with increasing sample depths in these cases. This is a strong indication that the growth of r_0 in standard volume-limited samples is caused by luminosity segregation, and a fractal explanation is disfavoured. Using a similar construction [3] found a growing r_0 in the Perseus Pisces survey (PPS). Our consistent results, both from the significantly larger SSRS2 and the CfA2 indicate that the PPS is too small in size to give reliable results. In the CfA2 [22] found also no significant growth of r_0 using a comparable method. [14] used mark correlation functions to quantify the luminosity dependency of the galaxy clustering in the SSRS2 in a scale dependent way, uninfluenced by inhomogeneities.

I limited my investigations to the question: what causes the growing amplitude of the two-point correlation function? Already with the current galaxy catalogues one is able to show that the amplitude of the two-point correlation function is depending on the luminosity of the galaxies. If one a priori assumes that the galaxy distribution is a fractal, these results may be interpreted as large fluctuations, which indeed are common in fractals. However, these fluctuations must conspire in all the samples considered here, to give the observed result of a constant amplitude of the two-point correlation function. Fluctuations in the morphological properties of the large scale distribution of galaxies have been detected out to a scale of $200h^{-1}\text{Mpc}$ [15]. However, these fluctuations are barely visible with two-point measures, and they are still compatible with fluctuations expected from a ΛCDM model. I did not comment on the topic, whether one already does see a turnover to homogeneity from current galaxy surveys [15], and whether $\xi(r)$ or $\Gamma(r)$ shows the more extended scaling regime. For a discussion see [23,5]. I expect that the completed Sloan Digital Sky Survey will offer conclusive evidence for these points.

* * *

I would like to thank Ruth Durrer and Francesco Sylos Labini for organising the workshop "Facts and Fiction in Cosmology" in April 2001 in Sils Maria. The discussions during this workshop generated the impulse to re-investigate these galaxy catalogues. For comments on an earlier version of the manuscript I am grateful to Alvaro Domínguez, Jose Gaite, Michael Joyce, Marco Montuori and especially to Claus Beisbart. I acknowledge support from the *Sonderforschungsbereich 375 für Astroteilchenphysik der DFG*.

REFERENCES

- [1] V.J. Martínez, E. Saar. *Statistics of the Galaxy Distribution*. Chapman and Hall, London, 2001.
- [2] P.J.E. Peebles. *The Large Scale Structure of the Universe*. Princeton University Press, Princeton, New Jersey, 1980.
- [3] F. Sylos Labini, M. Montuori, L. Pietronero. Scale invariance of galaxy clustering. *Physics Rep.*, 293:61, 1998.
- [4] L. Pietronero. The fractal structure of the Universe: Correlations of galaxies and clusters and the average mass density. *Phys. Rev. A*, 144:257, 1987.
- [5] J. Gaite, A. Domínguez, J. Perez-Mercader. The fractal distribution of galaxies and the transition to homogeneity. *ApJ*, 522:L5, 1999.
- [6] P.H. Coleman, L. Pietronero. The fractal structure of the Universe. *Physics Rep.*, 213:311, 1992.
- [7] M. Kerscher, I. Szapudi, A.S. Szalay. A comparison of estimators for the two-point correlation function. *ApJ*, 535:L13, 2000.
- [8] S.D. Landy, A.S. Szalay. Bias and variance of angular correlation functions. *ApJ*, 412:64, 1993.
- [9] L.N. da Costa, C.N.A. Willmer, P.S. Pellegrini, et al. The Southern Sky Redshift Survey. *AJ*, 116:1, 1998.
- [10] C. Benoist, S. Maurogordato, L.N. Da Costa, et al. Biasing in the galaxy distribution. *ApJ*, 472:452, 1996.
- [11] J.P. Huchra, M. Vogeley, M.J. Geller. The CfA redshift survey: Data for the South Galactic Cap. *ApJS*, 121:287, 1999.
- [12] W. Saunders, W.J. Sutherland, S.J. Maddox, et al. The PSCz catalogue. *MNRAS*, 317:55, 2000.
- [13] I. Szapudi, E. Branchini, C. Frenk, et al. The luminosity dependence of clustering and higher order correlations in the PSCz survey. *MNRAS*, 319:L45, 2000.
- [14] C. Beisbart, M. Kerscher. Luminosity and morphology dependent clustering of galaxies. *ApJ*, 545:6, 2000.
- [15] M. Kerscher, K. Mecke, J. Schmalzing, et al. Morphological fluctuations of large-scale structure: the PSCz survey. *A&A*, 373:1, 2001.
- [16] A.J.S. Hamilton. Evidence for biasing in the CfA survey. *ApJ*, 331:L59, 1988.
- [17] B.J.T. Jones, V.J. Martínez, E. Saar, J. Einasto. Multifractal description of the large-scale structure of the Universe. *ApJ*, 332:L1, 1988.
- [18] V.J. Martínez, M. Portilla, B.J.T. Jones, S. Paredes. The galaxy cluster correlation length. *A&A*, 280:5, 1993.
- [19] C.N.A. Willmer, L.N. da Costa, P.S. Pellegrini. Southern sky redshift survey: Clustering of local galaxies. *AJ*, 115:869, 1998.
- [20] P. Norberg, C.M. Baugh, Ed Hawkins, et al. The 2df galaxy redshift survey: The dependence of galaxy clustering on luminosity and spectral type. *MNRAS*, 332:827, 2002.
- [21] The SDSS Collaboration, I. Zehavi, M.R. Blanton, J.A. Frieman et al. Galaxy clustering in early Sloan Digital Sky Survey redshift data. *ApJ*, 571:172, 2002.
- [22] V.J. Martínez, B. Lopez-Martí, M.-J. Pons-Bordería. Does the galaxy correlation length increase with the sample depth. *Ap. J. Lett.*, 554:L5, 2001.
- [23] J.L. McCauley. The galaxy distributions: Homogeneous, fractal, or neither? *Fractals*, 6:109, 1998.